



**TARGET  
PHARMA**  
PHARMACEUTICAL

**MASTOGET**

**COMPOSITION:**

Each ml contains:

Drostanolone Propionate      100 mg  
Oil base                              q.s.

**Indications and usage:**

For use in females, for palliation of androgenresponsive recurrent mammary cancer in women who are more than one year but less than five years postmenopausal

**Contraindications:**

1. Known hypersensitivity to the drug
2. Males with carcinoma of the breast.
3. Males with known or suspected carcinoma of the prostate gland.
4. Women who are or who may become pregnant.
5. Patients with serious cardiac, hepatic or renal disease.

**Drug Interactions:**

Androgens may increase sensitivity to oral anticoagulants. Dosage of the anticoagulant may require reduction in order to maintain satisfactory therapeutic hypoprothrombinemia.

Concurrent administration of oxyphenbutazone and androgens may result in elevated serum levels of oxyphenbutazone In diabetic patients, the metabolic effects of androgens may decrease blood glucose and therefore, insulin requirements.

**Overdosage:**

Chronic ingestion of high doses of anabolic steroids can cause elevations in blood pressure, left ventricular hypertrophy and premature coronary artery disease.

**Precautions/Warnings:**

Women develop signs of virilism, with increased facial hair, male pattern baldness, acne, deepening of the voice, irregular menses and clitoral enlargement.

Changes in the larynx in women caused by anabolic steroids can result in a hoarse, deep voice.

The changes are irreversible.

Sodium and water retention can occur, and result in oedema; hypercalcaemia is also reported.

Androgen ingestion by a pregnant mother can cause virilization of a female fetus.

**Insulin resistance with a fall in:**

Glucose tolerance and hypercholesterolaemia with a fall in high density lipoprotein cholesterol have been reported.

**Dosage and administration:**

Intramuscular injection only usually 1.4 mg /kg of body weight. Thrice a week.

**How supplied**

Mastoget is supplied in 10x1 ml ampoules. Do not refrigerate or freeze. Do not store above 30° C & protect from light. Keep out of reach & sight of children. Do not use after the expiry date.

**Abuse in Sports**

Abuse of this product in sports is highly discouraged. Mastvolin does not take any responsibility for use of this product for any other purpose. This product should be used under proper medical supervision.

Effect of materials on plants or animals: This product may be harmful to contaminated plant and animal life

**Effect of chemicals on aquatic life:**

This product may be harmful to aquatic plant and animal life in contaminated bodies of water, especially it released in large quantities,